
	CRITERES DE SELECTION ETUDE NUTIDE 121	Identité patient (coller étiquette patient)
Version 1.0 du 30/11/2021	Investigateur en charge du patient : PI : Pr GHIRINGHELLI Mail : fghiringhelli@cgfl.fr <i>A contacter pour adresser/inclure patient externe au CGFL</i>	Arc : Céline SCHOUTIT Poste : 3427

« NUTIDE 121 »


Atezolizumab néoadjuvant et atezolizumab + bevacizumab en adjuvant après ablation percutanée par radiofréquence de CHC de petite taille : essai de phase II randomisé multicentrique.




VALIDATION DES CRITERES DE SELECTION

Critères d'inclusion :

Written informed consent and authorisation to use and disclose health information	<input type="checkbox"/> oui <input type="checkbox"/> non
	<input type="checkbox"/> oui <input type="checkbox"/> non

	CRITERES DE SELECTION ETUDE NUTIDE 121	Identité patient (coller étiquette patient)
Version 1.0 du 30/11/2021	Investigateur en charge du patient : PI : Pr GHIRINGHELLI Mail : fghiringhelli@cgfl.fr <i>A contacter pour adresser/inclure patient externe au CGFL</i>	Arc : Céline SCHOUTIT Poste : 3427


Ability to comprehend and willingness to comply with the requirements of this protocol, including the QoL questionnaires	
Female or male patients aged ≥ 18 years	<input type="checkbox"/> oui <input type="checkbox"/> non
Histologically- or cytologically-confirmed adenocarcinoma of the biliary tract (including gallbladder, intra and extra-hepatic biliary ducts and ampullary cancers) that is locally advanced, unresectable or metastatic (AJCC edition 8, 2018). Patients with measurable (as per RECIST v1.1 criteria) or non-measurable disease are permitted.	<input type="checkbox"/> oui <input type="checkbox"/> non
Life expectancy ≥ 16 weeks	<input type="checkbox"/> oui <input type="checkbox"/> non
ECOG performance status 0 or 1	<input type="checkbox"/> oui <input type="checkbox"/> non
Adequate biliary drainage with no evidence of ongoing infection. If applicable, treatable and clinically-relevant biliary duct obstruction has been relieved by internal endoscopic drainage/stenting at least 2 weeks previously or by palliative bypass surgery or percutaneous drainage prior to study entry, and the patient has no active or suspected uncontrolled infection. Patients fitted with a biliary stent should be clinically stable and free of signs of infection for ≥ 2 weeks prior to study entry. Patients with improving biliary function who meet all other inclusion criteria may be re-tested during the screening window.	<input type="checkbox"/> oui <input type="checkbox"/> non
1. Adequate bone marrow, hepatic, and renal function, as evidenced by: 2. • Absolute neutrophil count (ANC) $\geq 1,500/\mu\text{L}$ without colony-stimulating factor support 3. • Platelet count $\geq 100,000/\mu\text{L}$ 4. • Haemoglobin ≥ 9 g/dL without need for haematopoietic growth factor or transfusion support in prior 2 weeks	<input type="checkbox"/> oui <input type="checkbox"/> non

	CRITERES DE SELECTION ETUDE NUTIDE 121	Identité patient (coller étiquette patient)
Version 1.0 du 30/11/2021	Investigateur en charge du patient : PI : Pr GHIRINGHELLI Mail : fghiringhelli@cgfl.fr <i>A contacter pour adresser/inclure patient externe au CGFL</i>	Arc : Céline SCHOUTIT Poste : 3427


<p>5. • Total bilirubin $<2 \times$ upper limit of normal (ULN); does not apply to patients with Gilbert's syndrome. Consistent with inclusion criterion 7, patients whose whole bilirubin and biliary function is recovering may be re-tested during the screening period.</p> <p>6. • Alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST) $<5 \times$ ULN</p> <p>7. • Serum creatinine $\leq 1.5 \times$ ULN or creatinine clearance ≥ 45 mL/min actual or calculated by the Cockcroft-Gault method</p> <p>8. • International normalised ratio (INR) <1.5 and partial thromboplastin time (PTT) $<1.5 \times$ ULN; does not apply to patients on an anti-coagulant with stable dose 28 days prior to first dose.</p>	
<p>QTc interval <450 msec (males) or <470 msec (females), in the absence of bundle branch block. In the presence of bundle branch block with consequent QTc prolongation, patients may be enrolled based on a careful risk-benefit assessment.</p>	<input type="checkbox"/> oui <input type="checkbox"/> non
<p>Human Immunodeficiency Virus-infected patients who are healthy and have a low risk of Acquired Immunodeficiency Syndrome-related outcomes may be included in this study</p>	<input type="checkbox"/> oui <input type="checkbox"/> non
<p>Female patients of child-bearing potential (i.e., all women except those who are post-menopausal for ≥ 1 year or who have a history of hysterectomy or surgical sterilisation) must have a negative pregnancy test within 3 days prior to the first study drug administration. All patients of child-bearing potential must agree to practice true abstinence or to use two highly effective forms of contraception, one of which must be a barrier method of contraception, from the time of screening until 6 months after the last dose of study medication.</p>	<input type="checkbox"/> oui <input type="checkbox"/> non
<p>Male patients with a female partner must either have had a successful vasectomy or they and their female partner meet the criteria above (not of childbearing potential or practicing highly effective contraceptive methods).</p>	<input type="checkbox"/> oui <input type="checkbox"/> non

Critères de non inclusion :

<p>Combined or mixed hepatocellular/cholangiocarcinoma.</p>	<input type="checkbox"/> oui <input type="checkbox"/> non
<p>1. Prior systemic therapy for advanced or metastatic biliary tract cancer. However, prior chemotherapy in the adjuvant setting or low-dose chemotherapy given in conjunction with</p>	<input type="checkbox"/> oui <input type="checkbox"/> non

	CRITERES DE SELECTION ETUDE NUTIDE 121	Identité patient (coller étiquette patient)
Version 1.0 du 30/11/2021	Investigateur en charge du patient : PI : Pr GHIRINGHELLI Mail : fghiringhelli@cgfl.fr <i>A contacter pour adresser/inclure patient externe au CGFL</i>	Arc : Céline SCHOUTIT Poste : 3427

<p>radiotherapy in the adjuvant setting and completed at least 6 months prior to enrolment is permitted. The following prior interventions are allowed provided the patient has fully recovered:</p> <p>2. • Surgery: non-curative resection with macroscopic residual disease or palliative bypass surgery. Patients who have previously undergone curative surgery must now have evidence of non-resectable disease requiring systemic chemotherapy.</p> <ul style="list-style-type: none"> • Radiotherapy: prior radiotherapy (with or without radio-sensitising low-dose chemotherapy) for localised disease and there is now clear evidence of disease progression requiring systemic chemotherapy. • Photodynamic therapy: prior photodynamic therapy for localised disease with no evidence of metastatic disease or for localised disease to relieve biliary obstruction in the presence of metastatic disease provided there is now clear evidence of disease progression requiring systemic chemotherapy. <ul style="list-style-type: none"> • Palliative radiotherapy: palliative radiotherapy provided that all adverse events have resolved and the patient has measurable disease outside the field of radiation. 	
Prior treatment with or known hypersensitivity to NUC-1031, gemcitabine, cisplatin or other platinum-based agents or history of allergic reactions attributed to any parenteral excipients (e.g., dimethylacetamide [DMA], Cremophor EL, Polysorbate 80, Solutol HS 15).	<input type="checkbox"/> oui <input type="checkbox"/> non
Symptomatic central nervous system or leptomeningeal metastases.	<input type="checkbox"/> oui <input type="checkbox"/> non
History of other malignancies, except adequately treated non-melanoma skin cancer, curatively treated in situ cancer of the cervix, surgically excised or potentially curatively treated ductal carcinoma in situ of the breast, or low grade prostate cancer or patients after prostatectomy not requiring treatment. Patients with previous invasive cancers are eligible if treatment was completed more than 3 years prior to initiating the current study treatment, and the patient has had no evidence of recurrence since then	<input type="checkbox"/> oui <input type="checkbox"/> non
Concurrent serious (as deemed by the Investigator) medical conditions, including, but not limited to, New York Heart Association class III or IV congestive heart failure, history of congenital prolonged QT syndrome, uncontrolled infection, active hepatitis B or C, or other co-morbid conditions that in the opinion of the Investigator would impair study participation or cooperation.	<input type="checkbox"/> oui <input type="checkbox"/> non
Other acute or chronic medical, neurological, or psychiatric condition or laboratory abnormality that may increase the risk associated with study participation or investigational product	<input type="checkbox"/> oui <input type="checkbox"/> non

	CRITERES DE SELECTION ETUDE NUTIDE 121	Identité patient (coller étiquette patient)
Version 1.0 du 30/11/2021	Investigateur en charge du patient : PI : Pr GHIRINGHELLI Mail : fghiringhelli@cgfl.fr <i>A contacter pour adresser/inclure patient externe au CGFL</i>	Arc : Céline SCHOUTIT Poste : 3427

administration or may interfere with the interpretation of study results and, in the judgment of the investigator, would make the patient inappropriate for entry into this study.	
Prior exposure to another investigational agent within 28 days prior to randomisation	<input type="checkbox"/> oui <input type="checkbox"/> non
Major surgery within 28 days prior to randomisation; patient must have completely recovered from any prior surgical or other procedures.	<input type="checkbox"/> oui <input type="checkbox"/> non
Pregnant or breastfeeding	<input type="checkbox"/> oui <input type="checkbox"/> non
Residual toxicities from prior treatments or procedures which have not regressed to Grade ≤ 1 severity (CTCAE v5.0), except for alopecia or \leq Grade 1 peripheral neuropathy.	<input type="checkbox"/> oui <input type="checkbox"/> non
Concomitant use of drugs at doses known to cause clinically relevant prolongation of QT/QTc interval	<input type="checkbox"/> oui <input type="checkbox"/> non
Administration of a live vaccination within 28 days prior to randomisation.	<input type="checkbox"/> oui <input type="checkbox"/> non
Ongoing or recent (≤ 6 months) hepatorenal syndrome	<input type="checkbox"/> oui <input type="checkbox"/> non

Date : _____

Signature de l'investigateur : _____